



The Survey of Fixed Point Theory

Dr. Sunil G. Purane

Jamkhed Mahavidyalaya Jamkhed Dist: Ahmednagar (M.S.)

Corresponding Author- Dr. Sunil G. Purane

Email: sunilpuranemaths@gmail.com

DOI- 10.5281/zenodo.16758946

Abstract:

The theory of fixed point is one of the most important and powerful tools of the modern mathematics not only it is used on a daily bases in pure and applied mathematics but it is also solves a bridge between analysis and topology and provide a very fruitful are of interaction between the two.

Fixed Point Theory :

The theory of fixed points belongs to topology, a part of mathematics created at the end of nineteenth century. The famous French mathematician H. Poincare (1854-1912) was the founder of the fixed point approach. He had deep insight into its future importance for problems of mathematical analysis and celestial mechanics and took an active part in its development. Fixed point theory is a rich, interesting and exciting branch of mathematics. It is relatively young but fully developed area of research. Study of the existance of fixed points falls within several domains such as, classical analysis, functional analysis, operator theory, general and algebric topology. Fixed points and fixed point theorems have always been a major theoretical tool in fields as widely apart as topology, mathematical economics, game theory, approximation theory and initial and boundary value problems in ordinary and partial differential equations. Moreover, recently, the usefulness of this concept for applications increased enormously by the development of accurate and efficient techniques for computing fixed points, making fixed point methods a major tool in the arsenal of mathematics. Fixed point theory is equivalent to best approximation, variational inequality and the maximal elements in mathematical economics. The sequence of iterates of fixed point theory can be applied to find solution of a variational inequality and the best approximation theory.

The theory of fixed points is concerned with the conditions which guarantee that a map $T : X \rightarrow X$ of a topological space X into itself. admits one or more fixed points that is, points x in X for which $x = Tx$. For example, a translation, i.e. the mapping $T(x) = x + a$ for a fixed a , has no fixed point, a rotation of the plane has a single fixed point (the center of rotation), the mapping $x \rightarrow x^2$ of $\mathbb{R}$ into itself has two fixed points (0 and 1) and the projection of $\mathbb{R}^2$ into the x -axis has infinitely many fixed points (all points of the x -axis). Existence problems of the

type arise frequently in analysis. For example, the problem of solving the equation $p(z) = 0$, where p is a complex polynomial, is equivalent to find a fixed point of the self maps of $\mathbb{C}$. More generally, if T is any operator acting on a subset M of a linear space E , to show that the equation $Tu = u$ has a solution, is equivalent to show that the map T has a fixed point. The earliest fixed point theorem is that of Brouwer [11], who in (1912), proved that a continuous self mapping T of the closed unit ball has at least one fixed point, that is, a point x such that $Tx = x$. Several proof of this historic result can be found in the existing literature.

Another fundamental result after Brouwer's fixed point theorem was given by polish mathematicians S. Banach in (1922). Banach proved a theorem, which ensures under appropriate conditions, the existence and uniqueness of a fixed point. This result is popularly known as "Banach fixed point theorems" or the "Banach Contraction Principle". It states that a contraction mapping of a complete metric space into itself has a unique fixed point. It is the simplest and one of the most versatile results in fixed point theory. Being based on an iteration process, it can be implemented on a computer to find the fixed point of a contractive map, it produces approximations of any required accuracy. Due to its applications in various disciplines of mathematics and mathematical sciences, the Banach contraction principle has been extensively studied and generalized on many settings and fixed point theorems have been established.

There are a large class of mappings for which fixed point theorems have been studied. It includes contractive mappings, contraction of various order mappings, Ciric contraction, asymptotically regular, dencifying etc. Apart from single mappings, pair of mappings, sequence of mappings and family of mappings are also some of the classes of mappings that have interested mathematicians. There are many settings in which

the fixed point theorems have been studied. Metric spaces and its various generalizations, 2-metric spaces, uniform and quasi-uniform spaces, Banach spaces, normed spaces, 2-normed spaces, locally convex spaces, uniformly convex spaces etc. are some of the settings in which fixed point theorems have been proved and their useful applications have been found. Chu-Diaz [18] and Bryant [14] observed that it is sufficient for some iterates to be a contraction in order to get a unique fixed point. Rakotch [78] and Boyd-Wong [10] have attempted to generalize the Banach contraction principle by replacing the Lipschitz constant by some real valued function whose value is less than one. But generally, in order to accommodate a variety of continuous and discontinuous functions, attempts were made to replace the contractive conditions by some general form of mapping condition (called generalized contraction). In (1977), Rhoades [80] has made a comprehensive study and compared various contractive conditions which are scattered in the literature. He also introduced some new definitions. In (1978), Rhoades [81], enlarged his system, where a number of definitions were duplicated by Fisher work ([22], [23]). In (1980), Hedrus [32] defined the concept of generalized Banach contraction. In it appeared the diameter of a non-finite set in the inequality of definition, for the first time. This new definition was the base of many generalizations. These definitions were systematized by Park [68]. In 1992, Meszars [54] proved the equivalence of forty three contractive definitions and many inclusion relations. Recently, Rhoades [82] tried to obtain general results from which a lot of already known results follow as a corollary. He also obtained some new results. On the other hand, in (1976), Jungck [36] generalized the Banach contraction principle by introducing a contraction condition for a pair of commuting self mapping on metric space and pointed out the potential of commuting mappings for generalizing fixed point theorems in metric spaces [37]. Jungck's results have been further generalized by considering general type of contractive conditions on the pair of mappings by Das and Naik [19], Kasahara [44], Park ([68], [69], [70]), Ranganathan [79], Singh [95] and several others. Further generalizations have also been obtained by taking contractive type conditions for three self mappings on a metric space – one of the mappings commuting with other two by Qureshi and Awadhiya [76], Bhola and Sharma [8], Khan and Imdad [48], Fisher [26] etc.

In (1982) Sessa [88], initiated the tradition of improving commutativity conditions in metrical common fixed point theorems. While doing so Sessa [88] introduced the notion of weak commutativity. Motivated by Sessa [88], Jungck [38] defined the concept of compatibility of two mappings, which includes weakly commuting mappings as proper

subclass. After this definition there is a multitude of compatibility like conditions such as : compatibility of type (A) (Jungck, Murthy & Cho [39]), Compatibility of type (B) (Pathak and Khan [65]), compatibility of type (P) (Pathak et al. [66]), weak compatibility of type (A) (Lal, Murthy & Cho [52]), p-weak compatibility (Ume and Kim [110]) whose details can be seen in their introducing papers. In (1998), Jungck and Rhoades [40] termed a pair of mappings to be weakly compatible (or coincidentally commuting) if they commute at their coincidence point. In (2001), Ahmed and Rhoades [2] produced some common fixed point theorems for compatible mappings on complete metrically convex metric spaces thereafter in (2002) Aamri and Moutawaki [1] gave some new fixed point theorems under strict contractive conditions. In (2003) Som [108] obtained some common fixed point results for a weaker type of mappings than commuting or weakly commuting, called compatible mappings, satisfying a more general inequality condition. In (2004) Phaneendra [72] have obtained common fixed point theorems for a pair of self maps which commute at their coincidence points, called weakly compatible maps using the idea of an orbit relative to self maps.

It has been known since the paper of Kannan [42] that there exists maps possessing discontinuity in their domain but still admitting fixed points. However, in every case the maps involved were continuous at the fixed point. Recently some authors attempted to relax continuity requirement in such results and for the work of this kind one may refer to Pant ([61], [62], [63]), Singh and Mishra [103] and Pant, Lohari and Jha [64]. In (2001) Beg [5] proved an iteration scheme for asymptotically non expansive mappings on convex metric spaces. In (2003) he [6] obtained an iteration process for non process for non linear mappings in uniformly convex linear metric spaces. He also proved in [7] fixed point set function of non expansive random mapping on metric spaces. In (2003) Fisher & Duran [25] proved some fixed point theorems for multivalued mappings or orbitally complete uniform spaces. In (2004), Suzuki [109] generalized the result of Kannan [43]. In (2005), Popa [73] has improved the result of several author by removing the assumption of continuity, relaxing compatibility to the weak compatibility property and replacing the completeness of the spaces with a set of four alternative condition for four functions satisfying and implicit relations. In (2006) Proinov [75] established the Meir-Keeler type contractive conditions and the contractive definitions involving gauge functions. The concept of 2-metric spaces has been investigated initially by Gähler [27]. This concept was subsequently enhanced by Gähler ([28], [29]), White [112] and several others. On the other hand Iseki [33], Iseki-Sharma-Sharma [34], Khan-Fisher [47], Khan [46], Singh-Tiwari-Gupta [98]

and a number of other authors have studied the aspects of fixed point theory in the setting of 2-metric space.

In the last three decades, a number of authors Cho-Khan-Singh [17], Iseki [33], Iseki-Sharma-Sharma [34], Ram [77], Sharma ([91], [92]), Singh [(96), (97)], Singh-Ram (99) and Singh and Virendra (100) have studied the aspects of fixed point theory in the settings of 2-metric spaces. They have been motivated by various concepts already known for metric spaces and have thus introduced analogous of various concepts in the frame-work of the 2-metric space. Khan [45], Murthy-Chang-Cho-Sharma [56], Rhoades [83], Singh-Tiwari and Gupta [98] and Naidu-Prasad [57] introduced the concepts of weakly commuting pairs of self mappings, compatible pairs of self mappings of type (A) in a 2-metric spaces, and they have proved several fixed point theorems by using the weakly commuting pairs of self mappings, compatible pairs of self mappings of type (A) in a 2-metric spaces. In (2001), Naidu [58] has proved some fixed point theorems for pairs as well as quadruples of self maps on a 2-metric space satisfying certain generalized contraction condition. In (2001) B. Singh and R.K. Sharma [105] have proved some common fixed point theorems using the concept of compatible mappings in 2-metric spaces.

Browder and Petrysyn [13] introduced the concept of asymptotically regular maps at a point in a metric space which is defined as:

“A mapping of a metric space (X, d) into itself is said to be asymptotically regular at a point x in X if ”.

Using this concept, many authors proved various result on fixed point and common fixed points for such mappings in complete metric spaces, for the work of this kind one can be referred to Anderson et al [4], Guay-Singh [31], Sharma-Yuel [93]. Rhoades et. al. [84] introduced the concept of relative asymptotic regularity for a pair of mapping on a metric space and Jungck [38] proposed the concept of compatible mappings and weakly commuting mappings. Sessa [88] and others used both cited concepts and gave many interesting results.

Singh-Virendra [101] have proved a common fixed point theorem for three weakly commuting mappings by using the concept of relative asymptotic regularity of a sequence in 2-metric spaces. Later on using the idea of compatible mappings Singh and Sharma [105] generalized, the result of Singh and Virendra [101]. B. Singh, Chauhan and Sharma [104] extends the results of Iseki and others [35] for four compatible maps and proved a common fixed point theorem using the concept of relative asymptotic regularity of sequence. Nesic [60] give a general result about fixed points for asymptotic regular mappings on a complete metric spaces.

Asymptotic fixed point theory involves assumptions about the iterates of the mapping in question. It has a long history in non linear functional analysis and in fact the concept of ‘asymptotic contractions’ is suggested by one of the earliest version of Banach’s principle attributed to Caccioppoli [15]. In (2003), Kirik [51] introduced an asymptotic version of Boyd – Wong [10] result. In the literature of fixed point theory many authors have extensively studied. Common fixed point theorems in Banach spaces. Bose [9] proved same results by taking the domain a closed convex subset of a uniformly convex Banach spaces. Prasad [74], Sahani and Bose [85] extended the results of Bose [9], by relaxing the condition of convexity from the domain. Recently, Som [107] obtained some fixed point theorems in uniformly convex Banach spaces, which generalize some results of Prasad [74] and Sahani – Bose [85] with respect to their mappings and inequality conditions. In (1992) Ahmad and Imdad [3] obtained some common fixed point theorem for compatible asymptotic regular mappings in Banach spaces and generalized some known results with respect to their mappings and inequality conditions. In (2003), Penot [71] proves some non expansive mappings from a closed convex subset of uniformly convex Banach spaces into itself under some asymptotic contraction assumptions.

B. Gregus [30], proved a fixed point theorem in Banach space, which is called Gregus fixed point theorem and then many authors have obtained some fixed point theorems of Gregus type. (Fisher and Sessa [24]), Jungck [41], Mukherji and Verma [55] and many others. In (2001) Sushil Sharma and Bhavna Deshpande [94] have prove some fixed point theorem of Gregus type for compatible mapping in Banach Spaces.

Khan and Imdad [49] obtained some results on fixed point of certain involution in Banach spaces. In (2003), he has introduced the concept of composite involution in Banach spaces. Several interesting results using fixed point theory are given in approximation theory. During the last 130 years or so this area has attracted the attention of several mathematicians. An excellent reference is Cheney [16]. For a survey paper one is referred to M.L. Singh [102] afterwards the theorem of Brosowski [12] has been a basic important results which was the generalization of Meir-Keeler [53].

In (1979) Singh [106] improved the results of Brosowski [12], using a fixed point theorem of Jungck [36], Sahab, Khan and Sessa [87] generalized the results of Singh [106]. Pathak, Cho and Kang (67) gave an application of Jungck’s (41) fixed point theorem to best approximation theory. They extended the results of Singh (106) and Sahab et. al. (87). In (2001) Dhange [20] generalized the result of Sahab et. al. (87) in best approximation

theory under some weaker conditions. In (2004), Nashine [59] proved some fixed point theorem without star shaped ness condition of domain and linearity condition of mappings in setup domain and linearity condition of mapping in setup of normed linear space. Recently Vijayraju and Marudai [111] proved some results on common fixed compact mappings are established in the setting of normed linear space which is an extension of results of Sahab, Khan [86] and Dotson [21] as a consequence some applications of best approximations are established.

References

1. M. Aamri and D.E.I. Moutawakil., "Some new common fixed point theorem under strict contractive conditions", J. Math. Anal. Appl., 270 (2002), 187-188.
2. M.A. Ahmed and B.E. Rhoades., "Some common fixed point theorems for compatible mappings", Ind. J. pure appl. Math., 32 (8), (2001), 1247 – 1254.
3. Aqueel Ahmed and M. Imdad., "Relative asymptotic regularity and fixed point theorems", The Aligarh, Bull. Math., Vol. 14 (1992).
4. D.F. Anderson, M.D. Guay and K.L. Singh., "Fixed and Common Points in Convex Metric Spaces", Jnanabha., 18 (1988), 31-43.
5. Beg., "An iteration scheme for asymptotically nonexpansive mappings on Uniformly Convex metric spaces", Non linear Analysis forum., 6 (1) (2001), 27 – 34.
6. Beg., "An iteration process for non linear mappings in Uniformly Convex Linear Metric Spaces", Czezechoslovak. Math. J., 53 (2) (2003), 405 – 412.
7. Beg., "Fixed Point Set Function of nonexpansive random mapping on metric Spaces," Math. Inequalities Appl., 6 (3) (2003), 545 – 552.
8. P.K. Bhola and P.L. Sharma., "Common fixed point theorem for three maps", Bull. Cal. Math. Soc., 83 (1991), 398-400.
9. S.C. Bose., "Common Fixed Point of mapping in a Uniformly convex Banach Sapce", J. Lond. Math. Soc., 18 (1978), 151.
10. D.W. Boyd and J.S.W. Wong., "On non-linear contractions", Proc. Amer. Math. Soc., 20 (1969), 458 – 464.
11. L.E.J. Brouwer., "Uber Abbildungen Von Mannigfaltigke item," Math. Ann., 71 (1912), 97 – 115.
12. B. Brosowski., "Fixpunktsatze in der approximation theorie," Mathematic (Cluj), 11 (1969), 195.
13. F.E. Browder and W.V. Petryshyn., "The Solution by iteration non-linear functional equations in Banach Spaces", Bull. Amer. Math. Soc., 72 (1966), 571 – 575.
14. V.W. Bryant., "Uber Abbildungen Von Mannigfaltigke item", Math. Ann., 71 (1912), 97-115.
15. R. Caccioppoli., "Una teorana general Sull", esistenza di elemerrti uniti in una transfor mazione funzionale", Rend. Accad. Naz. Lincei., 11 (1930) 794 – 799.
16. E.W. Cheney., "Applications of fixed point theory to approximation theory", Proc. Approximation Theory and Application., Academic Press I, Edlaw and Sahnay (1976).
17. Y.S. Cho", M.S. Khan and S.L. Singh., "Common fixed points of weakly commuting mappings., "Univ U. Novam, Zb. Rad. Prirod. Mat. Fak. Ser. Mat., 18 (1) (1988), 129 – 142.
18. S.C. Chu and J.B. Diaz., "Remarks on a generalization of Banach Contraction mappings", J. Math. Analysis & Appl., 11 (1965), 440 – 446.
19. K.M. Das and K.V. Naik., "Common fixed point theorems for Commuting maps on a metric Space", Proc. Amer. Math. Soc., 77 (1979), 369-373.
20. B.C. Dhange., "A result in best approximation theory, 32 (8) 2001 , 1209 – 1216.
21. W.J. (J.) Dotson., "Fixed point theorems for non expansive mappings on starshaped Subsets of banach Spaces", J. London. Math. Soc., 4 (1972), 408 – 410.
22. B. Fisher., "Results on fixed points", Bull, Acad, Pol. Des. Sci., 25 (1977), 1253 – 1256.
23. B. Fisher., "Results on Common fixed points", Math. Japonica., 22 (1977), 335 – 338.
24. B. Fisher and S. Sessa., "On a fixed point theorem of Gregus," Internat, J. Math and Math. Sci., 9 (1986), 23.
25. B. Fisher and D. Turkoglu., "Fixed point of multi valued mappings in uniform spaces", Proc. Indian. Acad. Sci. (Math. Sci.), 113 (2) (2003), 183-187.
26. B. Fisher., "Three mappings with a common fixed point," Math. Sem. Notes., 10 (1982), 293 – 302.
27. S. Gahler., "2-metrische Raume and Ihre topologische strukture", Math. Machr., 26 (1963), 115 – 148.
28. S. Gahler., "Uber die uniformisierbarkeit 2 – metrischer raume", Math. Nachr., 26 (1963), 115 – 148.
29. S. Gahler., "Zur Geometric 2-metrischer raume", Rev. Roumaine Math. Pures. Appl., 11 (1966), 665 – 667.
30. M. Gregus., "A fixed point theorem in Banach Space," Boll. Um. Math. Ital.; (5) (17A) (1980), 198.
31. M.D. Guay and K.L. Singh, "Convergence of Sequence of iterates for a pair of mappings," J.Math. Phys. Sci., 18 (1984), 461-472.

32. M. Hedemus., "Some extensions of fixed point theorems", Publ. Inst. Math., 27 (41) (1980), 77.
33. K. Iseki., "Fixed point theorem in 2-metric spaces", Math. Sem. Notes Kobe. Univ., 3 (1975), 133-136.
34. K. Iseki, P.L. Sharma and B.K. Sharma., "Contraction type mappings on 2-metric spaces", Math. Japonica., 21 (1976), 67-70.
35. K. Iseki, S. Rajput and P.L. Sharma., "An extension of Banach Contraction principle through rational expressions", Math. Same Notes., 10 (1982), 677.
36. G. Jungck., "Commuting mappings and fixed points", Amer. Math. Monthly., 83 (1976), 261-263.
37. G. Jungck., "Periodic and fixed points and Commuting mappings", Proc. Amer. Math. Soc., 76 (1979), 333-338.
38. G. Jungck., "Compatible mappings and Common fixed points", Internat. J. Math. and Math. Sci., 2 (1986), 285-288.
39. G. Jungck, P.P. Murthy and Y.J. Cho., "Compatible mappings of type (A) and Common fixed point theorems," Math. Japan., 38 (1993), 381-390.
40. G. Jungck and B.E. Rhoades., "Fixed points for set – valued function without continuity", Indian. J. Pure Appl. Math., 29 (3) (1998), 227-238.
41. G. Jungck, "On a fixed point theorem of fisher and Sessa", Internat. J. Math and Math. Sci., 13 (1990), 497.
42. R. Kannan., "Some results on fixed points", Bull, Cal. Math. Soc., 60 (1968), 71.
43. R. Kannan., "Some results on fixed points II," Amer. Math. Monthly., 76 (1969), 405-408.
44. S. Kasahara., "On Some results on fixed points", Math. Sem. Notes., 6 (1978), 373-382.
45. M.S. Khan, "On Convergence of Sequence of fixed points in 2-metric space", Ind. J. Pure. Appl. Math, 10 (1979), 1062-1067.
46. M.S. Khan., "On fixed point theorem in 2-metric space", Publ. Inst. Math. (Beograd) (N.S.), 41 (1980), 107-112.
47. M.S. Khan and B. Fisher, "Some fixed point theorems for commuting mapping", Math. Nachr., 106 (1982), 323-326.
48. M.S. Khan and M. Imdad., "Some Common fixed point theorems", Glas. Math., 18 (38) (1983), 321-326.
49. M.S. Khan and M. Imdad., "Fixed points of certain involutions in Banach Spaces, J. Austral. Math. Soc., 37 (1984), 169-177.
50. Q.H. Khan and M. Imdad, "Remarks on fixed and coincidence points of Certain composite Involutions in Banach Spaces", Mathematical student., Vol-72, No. – 1-4 (2003), 165-169.
51. W.A. Kirik., "Fixed points of asymptotic contractions", J. Math. Anal. Appl. 277 (2003), 645-650.
52. S.N. Lal, P.P. Murthy and Y.J. Cho., "An extension of TELCI, TAS and fisher's theorem," J. Korean. Math. Soc., 33 (1996), 891-908.
53. G. Meinardus., "In variate bei Linearen Approximation", Arch. Rational. Mech. Anal., 14 (1963), 301-303.
54. J. Meszaros., "A comparison of various definitions of contractive type mappings", Bull. Cal. Math. Soc., 84 (1992), 167-194.
55. R.N. Mukherjee and V. Verma., "A note on a fixed point theorem of Gergus", Math. Japonica. 33 (1988), 745.
56. P.P. Murthy, S.S. Chang, Y.J. Cho and B.K. Sharma., "Compatible mappings of type (A) and common fixed point theorem", Kyungpook. Math. J., 32 (2) (1992) 203-216.
57. S.V.R. Naidu and J.R. Prasad., "Fixed point theorems in 2- metric spaces", Ind. J. Pure and Appl. Math., 17 (1986), 974-993.
58. S. Venkataratnam Naidu., "Some fixed point theorems in 2-metric spaces", Ind. J. Math & Math. Sci., 28 (11) (2001), 625-636.
59. Hemant Kumar Nashne., "Best approximation for compatible mappings in non-convex set", Ind. J. Math., Vol-46, No.2-3 (2004), 245-258.
60. Slobodan C. Nasic., "Result out fixed point of asymptotically regular mappings", Ind. J. Pure. appl. Math., 30 (5) (1999), 491-494.
61. R.P. Pant., "Common fixed point theorem for contractive mappings," J. Math. Anal. Appl., 226 (1998), 251.
62. R.P. Pant., "A Common Fixed point theorem under a new conditions", Ind. J. Pure. Appl. Math., 30 (1999).
63. R.P. Pant, "On Continuity at fixed points," J. Indian. Math. Soc., 67 (2000), 72 (1-5).
64. R.P. Pant, A.B. Lohari and K. Jha., "Meir – Keeler type fixed point theorem and reciprocal continuity", Bull. Cal. Math. Soc., 94 (6) (2002), 459-466.
65. H.K. Pathak, M.S. Khan., "Compatible mapping of type (B) and Common fixed point theorems of Gregus type," Czech. Math. Jour., 45 (120) (1995), 685-698.
66. H.K. Pathak, Y.J. Cho, S.S. Chang, and S.M. Kang., "Compatible mappings of type (P) and fixed point theorem in metric spaces and probabilistic metric spaces, Novi Sad. J. Math., 26 (2) (1996), 87-109.
67. H.K. Pathak, Y.J. Cho and S.M. Kang., "An applications of fixed point theorems." Internat. J. Math. and Math. Sci., 21 (1998), 467.
68. S. Park., "On general contractive type conditions", J. Korean. Math. Soc., 17 (1980), 131-140.

69. S. Park., "Fixed points of f-contractive maps", Rocky Mountain. J.Math., 8 (1978), 743-750.
70. S. Park., "An extension of fixed point theorem of Kasahara", Math. Sem. Notes, 7 (1979), 85-89.
71. Jean – Paul – Penot., "A fixed point theorems for asymptotically contractive mappings", Proc. Amer. Math. Soc., Vol (13), 8 (2003), 2371-2377.
72. T. Phaneendra., "Coincidence points of two weakly compatible self maps and Common fixed point theorem through orbits", Indian, J. Math., Vol. (46) (2004), No – 2 – 3, 173-180.
73. Valeriu Popa., "A Generalization of Mier-Keeler type common fixed point theorem for four non continuous mappings", Sarjevo. J. of Math., Vol. (1) (13) (2005), 135-142.
74. D. Prasad., Common fixed points of mappings in a uniformly Convex Banach Spaces with a New Functional inequality", Ind. J. Pure. appl. Math., 15 (2) (1984), 115.
75. Petko. D. Prinov., "Fixed point theorem in metric spaces", Non Linear Analysis., 64 (2006), 546-557.
76. K. Qureshi and R.K. Awadhiya., "Fixed point theorem for three mappings in Complete metric space," Math., Edu., 29 (3) (1995), 164-166.
77. B. Ram., "Existence of fixed points in 2-metric spaces", Indian. J. Pure and Appl. Math., 17 (1986), 974-993.
78. E.Rakotch., "A note on contractive mappings", Proc. Amer. Math. Soc., 13 (1968), 459-465.
79. S. Ranganathan., "Fixed points theorems for commuting mapping", Math. Sem. Notes., 6 (1978), 351-357.
80. B.E. Rhoades., "A comparison of various definitions of contractive mappings", Trans. Amer. Math. Soc., 226 (1977), 257-290.
81. B.E. Rhoades., "A collection of contractive definitions", Math. Sem. Notes., 6 (1978), 229-235.
82. B.E. Rhoades., "Proving fixed point theorems using General principle", Ind. J. Pure. Appl. Math., 27 (8) (1996), 741-770.
83. B.E. Rhoades., "Contraction type mappings an a 2-Metric space ", Math .Nachr., 91 (1979), 151-154.
84. B.E. Rhoades, S. Sessa, M.S. Khan, M. D. Khan., "Some fixed points in 2-metric Space", Math. Japonica., 29, No-4 (1984), 519-525.
85. D. Sahani and R. K. Bose., "Common fixed points of mappings in a uniformly convex banach space", Indian J. Pure and appl. Math., 15 (1984), 625.
86. S. A. Sahab and M. S. Khan., "Some results on best approximations", Univ. U. Novom. Sadu. Zb. Prirod –Mat. Fak. Ser. Mat., 17 (1987), 143-152.
87. S.A. Sahab, M. S. Khan and S. Sessa., "A result in best approximation theory", J. Approximations theory., 55 (1988), 349.
88. S. Sessa., "On a weak commutativity condition of mappings in fixed point considerations", Publ. Inst. Math. (Beograd), 32 (1982), 149-153.
89. A.K. Sharma., "On fixed points in 2-metric Spaces", Math. Seminar. Notes. Kobe Univ., (1979) 6, 467-473
90. A.K. Sharma., "A study of fixed points of mappings in metric and 2-metric spaces", Ph.D. Thesis (1979), Delhi Univ.
91. K. Sharma., "A generalization of Banach contractions principle in 2-metric spaces", Math. Seminar Notes. Kobe. Univ., 7 (1979), 291-292.
92. K. Sharma., "A note on fixed points in 2-metric spaces", Indian J. Pure and Appl. Math., 11 (1980), 1580-1583.
93. P. L. Sharma and A. K. Yuel., "Fixed point theorems under asymptotic regularity at a point", Math. Sem. Note., 10 (1984), 183-190.
94. Sushil Sharma and Bhavna Deshpande., "Fixed point theorem and its applications to best approximation theory", Bull. Cal. Math. Soc., 93 (2) (2001), 155-166.
95. S. L. Singh., "On common fixed points of commuting mappings", Math. Sem. Notes., 5 (1977), 131-134.
96. S. L. Singh., "Some Contraction type principles on 2-metric spaces and applications", Math. Seminar. Notes, Kobe. Univ., 7 (1979), 1-11.
97. S. L. Singh., "A fixed point theorem in 2-metric spaces", Math. Edu (Swain), 14 (1980), 53-54.
98. S. L. Singh, B.M.L. Tiwari and V. K. Gupta., "Common fixed points of commuting mappings in 2-metric spaces and applications", Math. Nachr., 95 (1980), 293-297.
99. S. L. Singh and B. Ram., "Note on the convergence of Sequence of mappings and their fixed points in a 2-metric space", Math. Seminar. Notes, Kobe Univ., 9 (1981), 181-185.
100. S. L. Singh and Virendra., "Coincidence theorems on 2-metric spaces", Indian J. Phy. Nat. Sci., 2 (1982), 32-35.
101. S. L. Singh and Virendra., "Relative asymptotic regularity and fixed points", Ind. Jour of Math., 31 (1989), 99-104.
102. M. L. Singh., "Applications of fixed point to approximation theory", Proc. Approximation theory and Application, (1985) Pitman.
103. S. L. Singh and S. N. Mishra., "Remarks on Jachymski's fixed point theorem for compatible maps", Indian J. Pure. Appl. Math., 28 (5), (1997), 611.
104. Singh, M. S. Chauhan and R. K. Sharma., "Relative asymptotic regularity and common

- fixed points”, Bull. Cal. Math. Soc., 89 (1997), 323-326.
105. Singh and R. K. Sharma., “Compatible mappings and fixed points” , Bull. All. Math. Soc., Vol. 16 (2001), 115-119.
106. S.P. Singh., “An application of a fixed point theorem to approximation theory”, J. Approximation theory., 25 (1979), 89.
107. Tanmoy. Som., “Common fixed point theorem for asymptotically commuting mappings in uniformly convex banach spaces”, Indian. J. Math., 31 (2) (1989),193.
108. T. Som., “Few common fixed point results for compatible mappings,” Bull. Cal. Math. Soc., 95 (4) (2003), 307-312.
109. Tomonari Suzuki., “Several fixed point theorems concerning ”, Fixed point theory and application., 3 (2004), 195-209
110. J. S. Ume and T. H. Kim., “Common fixed point theorems for weak compatible mappings”, Ind. J. Pure. Appl. Math., 32 (4) (2001), 565-571.