



Cypermethrin-exposed freshwater crab *Paratelphusa Jacquimontii* (Rathbun) and its toxic effects on protein in its Muscles and Gills

Dr. S. K. Parate

Assistant Professor, Dept. of Zoology

B.B. Arts, N.B. Commerce and B.P. Science College, Digras

Corresponding Author – Dr. S. K. Parate

DOI - 10.5281/zenodo.15259644

Abstract:

Paratelphusa Jacquimontii (Rathbun) were exposed the concentration of cypermethrin i.e. 0.025 ml/L. Total protein contents showed decreasing trend over the period of four days.

Keywords: *Paratelphusa Jacquimontii*, *Cypermethrin* (Pyrethroid) Muscle, Gills, Total Protein

Introduction:

Pyrethroid are potential insecticides derived from the flower heads of *Chrysanthemum cinerariifolium* species [1]. Pyrethroid insecticides are widely used in the agricultural operation to control pest. Musa Galadima et.al, have reported that pyrethroid has harmful effect on the fish population and also direct effects on the human beings [2]. Particularly very little information exists regarding the influence of pyrethroids on the various systems of the crab. A new class of agricultural insecticides, the synthetic pyrethroid have become popular for their high toxicity to the wide range of the insects including resistant strains and low toxicity to mammals and birds [3]. The correspondence between aqueous concentration and LC₅₀ for test animals, demonstrated the importance for the quantifying the bio-available portion of the pyrethroid in the field samples to

characterize accurately the environmental risk associated with pyrethroid runoff after agricultural applications [4] [5].

Cypermethrin is one of the recent pyrethroids being used for the cotton crop pest in India [6]. Higher concentration of the toxicants in aquatic environment causes adverse effect on the non-target aquatic organism. Nominal sediment concentrations of cypermethrin are of 0.1 mg/kg (static) or 0.1 mg/kg (flow through) resulted in the mortality in pink shrimp (*Penaeus duoraum*) in the static and flow-through the test system [7].

Present study was undertaken to investigate the effect at biochemical level of a non-target species. Changes involved in total protein profile in the muscles and gills of the crab, *Paratelphusa Jacquimontii*, exposed to the pyrethroid pesticide cypermethrin at different experimental conditions was studied.

Table 1: total protein content in crab, *P. Jacquimontii* (Rathbun) under different experimental condition is for 4 days.

Tissue	N(N)	EA (N)	EAI +N	CP + CP	EA CP	EAI + CP
Muscles	6.025 + 0.25	6.516 +	6.492 +	6.437 +	6.384	5.412 ** +
		0.27	0.31	0.27	+0.28	0.27
Gills	4.021+0.02	4.011	4.092	4.088	3.074	3.068
		+0.21	+0.31	+0.02	+0.24	+0.22

(All values are significant at * p<0.05 and **p <0.01)

Where N: Normal crab, EA : Eyestalk ablated crab, EAI: Eyestalk ablated crab after administration of eyestalk extract injection, CP: Cypermethrin

Martial and Methods:

The crab *Paratelphusa Jacquimontii* (Rathbun) were collected as local fish market from Digras city, Vidarbha, Maharashtra State. The crabs were acclimatized to the laboratory condition for 7 days prior to the experimentation. Healthy active, adult medium sized, living crabs were selected for the experimental. Based on the predetermined LC₅₀ values pesticide concentration was prepared for normal, eyestalk ablated and eyestalk ablated but extract injected batches of crabs. Experimental animals were scarified after experimental period and muscles as well as gills dried at 70⁰c to 80⁰c in an oven, till constant weights were obtained.

The dried powders of the tissues of the normal, eyestalk abled and eyestalk ablated and extract injected animals were used for the estimation of protein. The results of estimation were expressed as mg/gm dry weight of the tissues.

Result and Discussion:

Pesticides due to their potential toxicity produce biochemical changes in the tissues and organs of exposed animals. In the present study. Total protein content decreased in the gills and muscles due to cypermethrin stress. In crab, prominent total protein decrease was observed cypermethrin exposure to an increase in the acidic protein in the beginning, which was followed by decrease in the total proteins till 48 hrs. It is well established that the total proteins are used as an alternative source of energy especially under that stress conditions [8].

Similar observation was made by Srivastava et.al (2013) on freshwater crab, *Baraytelphusa guerini* exposed to cuprous oxide [9]. Deshpande et.al (2014) also recorded a decline in protein content in the

crab *barytelphusa guerini* after exposure to zinc sulphate and copper sulphate [10].

It may be concluded from the present study that the protein was utilized for the production of the energy to mitigate the pesticide stress. Such change in metabolic activity cause metabolic rearrangement in the living systems. There was decrease in the total protein content in the muscle and gills of crab during early hours of exposure to cypermethrin. Similar result is reported by several investigators. Further total protein content might have acted as a fatigue retardant due to the effect of pesticide on the body parts [11]. Total protein levels in muscle and gills showed a decrease during exposure period of 24, 48, 72 and 96 hrs.

Eyestalk ablation caused marked changes in total protein, nevertheless after eyestalk extract injection to ablated crabs the levels of the total protein was restored[12]. In muscles as well as gills the total protein contents decreased in the normal and eyestalk ablated, crab and again levels were restored after eyestalk extract was injected.

Injection of eyestalk extracts reestablishes low levels of proteins in the ablated animals. On injection of extract protein synthesis is again inhibited, resulting in the restoration of the original condition of low protein contents, thus the experiments of eyestalk ablation d injection of eyestalk extract in to ablated crabs provides good evidence demonstrating a protein synthesis inhibitory factor in the eyestalk of the crab. Further evidence comes from the observation that eyestalk removal increases the protein levels in the all the tissues [14].

References:

1. Hodoşan C, Gîrd CE, Ghica MV, Dinu-Pîrvu CE, Nistor L, Bărbuică IS, Marin ŞC, Mihalache A, Popa L. "Pyrethrins

- and Pyrethroids: A Comprehensive Review of Natural Occurring Compounds and Their Synthetic Derivatives”, *Plants (Basel)*. 2023 Nov 29;12(23):4022. doi: 10.3390/plants12234022. PMID: 38068657; PMCID: PMC10707950.
2. Musa Galadima, Simranjeet Singh, Akshay Pawar, Sutripto Khasnabis, Daljeet Singh Dhanjal, Amith G Anil, Pratistha Rai, Praveen C. Ramamurthy, Joginder Singh, "Toxicity, microbial degradation and analytical detection of pyrethroids: A review", *Environmental Advances*, Volume 5, 2021, 100105, ISSN 2666-7657, <https://doi.org/10.1016/j.envadv.2021.10.0105>.
 3. Madara Ranatunga, Claudette Kellar, Vincent Pettigrove, "Toxicological impacts of synthetic pyrethroids on non-target aquatic organisms: A review", *Environmental Advances*, Volume 12, 2023, 100388, ISSN 2666-7657, <https://doi.org/10.1016/j.envadv.2023.100388>.
 4. Saha, Suchismita & Kaviraj, Anilava. (2008). Acute Toxicity of Synthetic Pyrethroid Cypermethrin to Some Freshwater Organisms. *Bulletin of environmental contamination and toxicology*. 80. 49-52. 10.1007/s00128-007-9314-4.
 5. Lichun Li, Shugui Liu, Yi Yin, Guangming Zheng, Cheng Zhao, Lisha Ma, Qi Shan, Xiaoxin Dai, Linting Wei, Jiawei Lin, Wenping Xie, "The toxicokinetics and risk assessment of pyrethroids pesticide in tilapia (*Oreochromis mossambicus*) upon short-term water exposure", *Ecotoxicology and Environmental Safety*, Volume 241, 2022, 113751, ISSN 0147-6513, <https://doi.org/10.1016/j.ecoenv.2022.11.3751>.
 6. Ayaz Ahamad, Jitendra Kumar, "Pyrethroid pesticides: An overview on classification, toxicological assessment and monitoring", *Journal of Hazardous Materials Advances*, Volume 10, 2023, 100284, ISSN 2772-4166, <https://doi.org/10.1016/j.hazadv.2023.10.0284>.
 7. Willis, Kate & Ling, Nicholas. (2004). Toxicity of the aquaculture pesticide cypermethrin to planktonic marine copepods. *Aquaculture Research*. 35. 263 - 270. 10.1111/j.1365-2109.2004.01008.x.
 8. Xianyun Ren, Qiong Wang, Qiong Wang, Huixin Shao, Huixin Shao, Yao Xu, Yao Xu, Ping Liu, Ping Liu, Jian Li, Jian Li, "Effects of Low Temperature on Shrimp and Crab Physiology, Behavior, and Growth: A Review", *Front. Mar. Sci.*, 21 October 2021,
 9. *Sec. Aquatic Physiology*, Volume 8 - 2021 | <https://doi.org/10.3389/fmars.2021.746177>
 10. Srivastava, J. & Das, Shobha & Pandey, Dr Rakesh & Das, Vijai. (2013). Acute toxicity and behavioural responses of freshwater crab *Barytelphusa guerini* to chlorpyrifos exposure. *J Appl Biosci*. 39. 56-59.
 11. Deshpande, Manoj & Nagthane, Nagesh & Nagrale, Narayan & Ashwini, Jagtap & Barde, Ravi. (2014). Pesticides induces alterations in protein content of fresh water male crab, *Barytelphusa guerini* (h. Milne - edw... Editor Biolife Journal Pesticides induces alterations in protein content of fresh water male crab, *Barytelphusa guerini* (h. Milne - edwards), from Godavari basin, Nanded, Maharashtra.
 12. Parate S K, "Effect of Cypermethrin on Lipid Level of Freshwater Crab *P. Jacquimontii* Hepatopancreas and Muscle", *Bioscience Discovery*, 4(1):128-132, Jan. 2013 ISSN: 2229-3469.
 13. Shuang Liang, Chuigen Zeng, Jian Liang, Xuan Luo, Xiaolong Gao, Caihuan Ke, Yongren Li, Limei Chen, Huiru Liu, Yongjun Guo, "Effects of eyestalk ablation on growth, feeding behavior, movement and circadian gene expression in *Haliotis discus hannai*", *Aquaculture Reports*, Volume 33, 2023, 101807, ISSN 2352-5134, <https://doi.org/10.1016/j.aqrep.2023.101807>.